

# DUAL ACTION WEED CONTROL FOR TURFGRASS PROFESSIONALS

## **NEGATE**<sup>®</sup>

Negate<sup>®</sup> controls poa annua, onion grass, ryegrass, and tough to kill broadleaf weeds.









Ask your Quali-Pro agent about Negate<sup>®</sup> today!

Distributed by:



www.centaur-asiapacific.com info@centaur-asiapacific.com

\*Hong Kong, Macau, Singapore, Malaysia & Vietnam.

#### Quali-Pro<sup>®</sup> Negate<sup>®</sup>

GROUP B HERBICIDE

QUALI-PRO<sup>®</sup> NEGATE<sup>®</sup> is a unique dual powered post-emergent Sulfonylurea herbicide for grassy and broadleaf weed control in established warm season turfgrass. Contains Rimsulfuron 16.67%, Metsulfuron methyl 20%. Available in 110g. Sulfonylurea herbicides are effective at low use rates (1-5% of other herbicides). Controls Ryegrass, Poa annua and other cool season grasses as well as other tough-to-control Broadleaf weeds such as Clover, Dandelion, Henbit and many others.

#### **Other Takeways**

- · Caution S5 Schedule.
- Synergistic formulation which provides superior control at low rates compared to other leading products.
- Provides the broadest spectrum of control within the ALS inhibitor class of chemistry.
- Low odour formulation that's excellent for commercial and public applications.
- Labelled for use on both Couch and Zoysia turf.
- Outstanding over-seeded Couchgrass transition and Zoysiagrass maintenance with superior cost effective performance.

### Negate's Synergistic Performance

#### Shoot Absorption

- Herbicide covers the leaf surface of the target weed.
- The product becomes lipophilic and moves across the cell membrane, then is translocatedthroughout the plant.
- This disrupts the branch chain amino acids, which causes internal plant toxicity and ultimately death.
- NEGATE is truly systemic and moves throughout the plant.

#### **Root Absorption**

- Herbicide particles that are not absorbed by the foliage can also be absorbed by the plant's root system as particles move downward through the soil.
- As NEGATE is taken up by the roots, it is also translocated throughout the plant.

SITUATION	WEEDS CONTROLLED	RATE/Ha
Turf	Winter Grass (Poa annua)	110 g/ha in
	Ryegrass (Lolium perenne)	200-500 L
Common	Onion Grass	of water per hectare
Couch, Hybrid Couch, Zoysia, Qld Blue Couch	Broadleaf Weeds including; Bindii (Soliva sessilis), Caustic Weed (Chamaesyce prostrata), White Clover (Trifolium repens), Cobblers Peg (Bidens pilosa), Creeping Indigo (Indigofera spicata), Creeping Oxalis(Oxalis spp.), Plantain (Plantago lanceolata), Tridax Daisy (Tridax procumbens)	





Hong Kong | Singapore | Malaysia | Australia Email: info@centaur-asiapacific.com Website: www.centaur-asiapacific.com



™Trademarks or ®Registered trademarks of an ADAMA Agricultural Solutions Ltd Company. \*Registered trademarks. Please note: This information is not intended to replace the product label. Always read the complete product label appearing on the container before opening or using products. ADA20102